A Glossy Ibis (*Plegadis faleinellus*) seen by myself at Yangebup Lake during January and February, 1957, and by Miss J. Russell and B. V. Teague in March, was probably the same individual seen by C. F. H. Jenkins ( $W.A.\ Nat.$ , 6:55) at Bibra Lake in November, 1956.

—JULIAN R. FORD, Fremantle.

Yellow-billed Spooubill and White Ibis at Australind.—On April 5, 1958, my son, Lindsay, and I inspected a pool near the Wellesley River and adjacent to the Brunswick-Australind road. Ten large white birds were present and at first sight all appeared to be White Egret (Egretta alba). One bird, however, attracted immediate attention by feeding in a typical spoonbill manner, i.e., pivoting head, neck and body from side to side with bill deeply submerged in the water in which the bird was standing. After some little time the bird raised its head, revealing a typical "spoon" bill, yellow in colour. Within a minute or so, the bird placed its head behind its wing and apparently went to sleep.

Only then could we give our attention to another bird about three feet from the Spoonbill, of similar form and also with head behind wing. We would have judged this to be a second Spoonbill, but whereas the legs of the Spoonbill were plainly yellow, those of its companion were black. As both birds had apparently settled

down, we left the pool for a time.

Returning four hours later, we found all our white birds perched in a tree near the pool. All took to flight, a Yellow-billed Spoonbill (Platalea flavipes) and a White Ibis (Threskiornis molucca) separating from eight White Egret. The two former eventually settled in the pool while the Egret perched in a second tree nearby.

It seemed that the Spoonbill and the Ibis were fairly closely associated with each other and that both were loosely associated

with the Egrets.

The Ibis appeared immature, i.e. head and neck were apparently fully feathered. In this respect it resembled five White Ibis which I saw by an ox-bow of the Preston River at Pieton on May 15, 1957.

-ERIC H. SEDGWICK, Collie.

Little Eagle in Metropolitan Area.—A light phase Little Eagle (Hieraaëtus morphnoides) was seen gliding over the shore of Monger's Lake, Leederville, on June 4, 1956. It was identified from several characters, including the underwing pattern. The eagle was heekled for some time by a pair of Magpies and gave the plaintive whistling call usually given by this species when being heekled. Serventy (Birds of the Swan River District, 1948, p. 32) gives only two records for the district (Cannington and Pickering Brook), neither of which is close to the city.

-J. H. CALABY, Canberra.

A Yellow Silvereye at Mandurah.—Further to the note by A. Jones on a yellow mutant of the Western Silvereye (Zosterops australasiae) at Windy Harbour (W.A. Nat., 6 (5): 127) I have to report a similar observation. In January 1958, while fishing at the